



# Clubroot Factsheet

Series 1 for nurseries

sheet 6 of 7

Monitor for clubroot in the nursery.

## Fact:

Clubroot can be difficult to detect in the nursery, however regular routine sampling will provide peace of mind for you and your customers.

Recognising clubroot is the first step in disease prevention.

- *Examine* seedlings trays at random on a regular basis, increasing in frequency over the summer months.
- *Remove* 2-3 seedlings from each tray and examine roots for small galls or deformities. Small galls or nodules along the root hairs are indicative of clubroot disease.
- *Train* staff to recognise signs of the disease and report to management with their findings.



- A *diagnostic test* (see fact sheet 1) is now available to assist with confirmation of diagnoses.

A PUBLICATION OF THE NATIONAL CLUBROOT PROJECT. JUNE 2003.

Supported by DPI (VIC), Agriculture WA, DPIWE (TAS), DPI (QLD) and NSW Agriculture.

Disclaimer: This publication may be of assistance to you but the State of Victoria and its officers do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purposes and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on any information in this publication.

© State of Victoria, Department of Primary Industries 2003

This publication is copyright. No part may be reproduced by any process except in accordance with the provisions of the Copyright Act 1968.