

Vegetable *Matters-of-Facts*

Based on research funded by the Australian Asparagus Council,
Horticulture Australia Limited and Department of Primary Industries –Victoria



Number 8
October
2003



Asparagus

Stemphylium (Purple Spot) of Asparagus

Main controls:

- Chemical fungicides
- Crop scouting
- Weed control
- Seed treatment
- Disease forecasting
- Remove fern debris



What is "Stemphylium" (Purple Spot)

Stemphylium of asparagus (also known as Purple Spot) is caused by the fungus *Stemphylium vesicarium*. Purple Spot is a major worldwide disease of asparagus.

Losses from this disease are from spotting of spears which reduces marketability. Repeated defoliation of ferns can reduce yield in following crops. The disease has caused severe problems for Southern Victorian asparagus growers in recent years.

Typical symptoms

Infected **spears** have elliptical sunken, purplish spots with brown centres which lead to rejection of product.

Asparagus **fern** infected with Purple Spot has brown spots with dark purple margins.

Black **over-wintering** structures appear as the fern dies back and these are the main source of infection for newly developing spears in spring.

More about Purple Spot

- The primary source of the disease is from infected fern debris from the previous crop.
- Infection occurs through natural openings and wounds on the newly emerging spears.
- Purple Spot can be severe during a cool, wet growing season. Twelve continuous hours of leaf wetness are necessary for infection.
- Purple Spot over-winters on fern debris. In spring, spores released from last year's infected fern debris are spread by wind and water to the new spears.




“Strategies to control *Stemphylium* (Purple Spot) of Asparagus”

HAL Project: VX 01024

Results to date



Controlling *Stemphylium* (Purple Spot)



Current methods

The control of Purple Spot involves a combination of crop management practices and well-timed chemical treatments.

Chemical Control:

In trials, the fungicide chlorothalonil was effective in controlling Purple Spot on the fern. Other effective chemicals are needed to reduce the danger of the disease becoming resistant to chlorothalonil.

NOTE: Chlorothalonil is not registered for use on Asparagus

Disease Forecasting:

Purple Spot can occur whenever weather conditions are right. Twelve hours of leaf wetness are necessary for infection. Crop scouting for purple spot on green fern is recommended after prolonged wet conditions.

For new plantings, consider orienting the block so that wind can blow along the rows and dry the foliage quickly.

Removing Fern Debris:

Burying or burning asparagus debris will help reduce infectious material and air-borne spores, even if the debris does not decay before harvest.

However total removal of fern debris is difficult and it is likely that enough infectious material will survive to infect emerging spears.

Weed Control:

Volunteer asparagus seedlings can become infected during the harvest season and are a source of disease. Mechanically removing volunteer seedlings can help control the spread of disease.



Are you on our mailing list ?

If you would like to receive your own copy of *Vegetable Matters-of-Facts* or have changed your address, please mail or fax this coupon to:

Craig Murdoch, Private Bag 15, Ferntree Gully Delivery Centre 3156; Fax (03) 9800 3521.

Name: _____

Address: _____

Postcode _____ Email _____

Organisation/Business _____

Crops grown _____

Seed treatment

Stemphylium, *Fusarium* and *Botrytis* (disease-causing fungi) were detected on asparagus seed.

A simple treatment to consider is to soak seed for 10 minutes in 7 parts methylated spirit and 3 parts water. This will rid seed of most fungal diseases. Unfortunately, *Stemphylium* can also be an internal infection and cannot be adequately controlled by surface sterilisation.

Crown testing

Random tests showed that 4 out of 6 one-year-old crown lots were infected with *Stemphylium* and all were infected with *Fusarium* root rot.

Decay of fern debris

Asparagus fern is very slow to decay in cold dry soil. Other methods to accelerate decomposition need to be developed.

Fungicide trials

In trial conditions, only the fungicide chlorothalonil provided a level of protection from disease at the green fern stage.

NOTE: Chlorothalonil is not registered for use on Asparagus

Future work

- Test low-toxicity fungicides for use in disease management to avoid chemical resistance developing.
- Test use of hot lime to aid decomposition of fern debris.
- Can crop scouting help growers.
- Disease forecasting from weather data.

Contact

Gisele Irvine or Martin Mebalds (seed testing)
DPI-Knoxfield, (03) 9210 9222

Weblinks used to prepare this information :

<http://www.apsnet.org/online/feature/asparagus/top.htm>

<http://pearl.agcomm.okstate.edu/plantdiseases/f-7646.html>

<http://www.extension.umn.edu/distribution/horticulture/components/1861a.html>

<http://attra.ncat.org/attra-pub/PDF/asparagus.pdf>

<http://www.anrcatalog.ucdavis.edu/pdf/7234.pdf>

http://commserv.ucdavis.edu/CEImperial/Aspar_03.pdf

For more information please contact your local VegCheque officer.

Neville Fernando	Gippsland	5152 0616
Sally-Ann Henderson	Northern Vic	5051 4500
Bruce Fry	South West Vic	5233 5510
Craig Murdoch	Melbourne	9210 9222

Vegetable Matters-of-Facts is published as part of DPI's VegCheque extension program.

Editor: Peter Carr, DPI-Knoxfield, 03 9210 9222.

Disclaimer: This publication may be of assistance to you but the state of Victoria and its employees do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purpose and therefore disclaims all liability for any error, loss or other consequences which may arise from you relying on any information in this publication.

© The State of Victoria, Department of Primary Industries, 2003.

ISSN: 1445-5676